

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
14 March 2002 (14.03.2002)

PCT

(10) International Publication Number  
**WO 02/20519 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 417/12**,  
A61K 31/44, A61P 3/00

(74) Agent: **RUTTER, Keith**; SmithKline Beecham, Corporate Intellectual Property (CN9.25.1), 980 Great West Road, Brentford, Middlesex TW8 9GS (GB).

(21) International Application Number: PCT/GB01/03991

(22) International Filing Date:  
5 September 2001 (05.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0021865.1 6 September 2000 (06.09.2000) GB

(71) Applicant (for all designated States except US):  
**SMITHKLINE BEECHAM P.L.C.** [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **CRAIG, Andrew, Simon** [GB/GB]; Glaxo SmithKline, Old Powder Mills, Near Leigh, Tonbridge, Kent TN11 9AN (GB). **MILLAN, Michael** [AU/GB]; Glaxo SmithKline, Old Powder Mills, Near Leigh, Tonbridge, Kent TN11 9AN (GB).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



**WO 02/20519 A1**

(54) Title: THE HYDROCHLORIDE SALT OF 5-[4-[2-(N-METHYL-N-(2-PYRIDYL)AMINO)ETHOXY]BENZYL]THIAZOLIDINE-2,4-DIONE

(57) Abstract: A novel pharmaceutical compound 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione hydrochloride, a process for preparing such a compound, a pharmaceutical composition comprising such a compound and the use of such a compound in medicine.

## THE HYDROCHLORIDE SALT OF 5-[4-[2-(N-METHYL-N-(2-PYRIDYL)AMINO)ETHOXY]BENZYL]THIAZOLIDINE-2,4-DIONE

This invention relates to a novel pharmaceutical, to a process for the preparation of the pharmaceutical and to the use of the pharmaceutical in medicine.

5 European Patent Application, Publication Number 0,306,228 relates to certain thiazolidinedione derivatives disclosed as having hypoglycaemic and hypolipidaemic activity. The compound of example 30 of EP 0,306,228 is 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione (hereinafter also referred to as "Compound I").

10 International Patent Application, Publication Number WO94/05659 discloses certain salts of the compounds of EP 0,306,228 and in particular the maleic acid salt.

It has now been discovered that Compound I forms a novel hydrochloride salt (hereinafter also referred to as the "Hydrochloride") that is particularly stable and hence is suitable for bulk preparation and handling. The Hydrochloride also has a high melting  
15 point, shows particularly good aqueous solubility and possesses good bulk flow properties. The Hydrochloride is therefore surprisingly amenable to large scale pharmaceutical processing and especially to large scale milling.

The novel form can be prepared by an efficient, economic and reproducible process particularly suited to large-scale preparation.

20 The novel Hydrochloride also has useful pharmaceutical properties and in particular it is indicated to be useful for the treatment and/or prophylaxis of diabetes mellitus, conditions associated with diabetes mellitus and certain complications thereof.

Accordingly, the present invention provides 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione hydrochloride characterised in that  
25 it:

- (i) provides an infrared spectrum substantially in accordance with Figure 1; and/or
- (ii) a Raman spectrum substantially in accordance with Figure 2; and/or
- (iii) an X-Ray powder diffraction pattern (XRPD) substantially in accordance with Figure 3; and
- 30 (iv) a Solid State  $^{13}\text{C}$  NMR spectrum substantially in accordance with Figure 4.

In one favoured aspect, the Hydrochloride provides an infrared spectrum substantially in accordance with Figure 1.

In a further aspect, the Hydrochloride provides a Raman spectrum substantially in accordance with Figure 2.

35 In one favoured aspect, the Hydrochloride provides an X-Ray powder diffraction pattern (XRPD) substantially in accordance with Figure 3.

In one favoured aspect, the Hydrochloride provides a Solid State  $^{13}\text{C}$  NMR spectrum substantially in accordance with Figure 4.

The Hydrochloride also provides provides a melting point in the range of from 160 to 168°C, such as 163 to 167 °C, for example 167 °C

The present invention encompasses the Hydrochloride isolated in pure form or when admixed with other materials. Thus in one aspect there is provided the

5 Hydrochloride in isolated form.

In a further aspect there is provided the Hydrochloride in a purified form.

In yet a further aspect there is provided the Hydrochloride in crystalline form.

Also, the invention provides the Hydrochloride in a solid pharmaceutically acceptable form, such as a solid dosage form, especially when adapted for oral  
10 administration.

Moreover, the invention also provides the Hydrochloride in a pharmaceutically acceptable form, especially in bulk form, such form being particularly capable of being milled. The invention therefor also provides the the Hydrochloride in a milled form.

Furthermore, the invention provides the Hydrochloride in a pharmaceutically  
15 acceptable form, especially in bulk form, such form having good flow properties, especially good bulk flow properties.

The invention also provides a process for preparing the Hydrochloride, characterised in that 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione (Compound I) or salt thereof, preferably dispersed or dissolved in a suitable  
20 solvent, is reacted with a source of chloride ion; and thereafter the Hydrochloride is recovered.

Suitably, Compound (I) is used in the reaction, especially in a protonated form.

Suitably, a salt of Compound (I) is used in the reaction..

Compound (I) is preferably present in a protonated form.

25 A suitable solvent is an alkanol, for example propan-2-ol, or a hydrocarbon, such as toluene, a ketone, such as acetone, an ester, such as ethyl acetate, an ether such as tetrahydrofuran, a nitrile such as acetonitrile, a halogenated hydrocarbon such as dichloromethane, an organic acid such as acetic acid, or water, or a mixture thereof.

Conveniently, the source of chloride ion is a solution of hydrogen chloride in an  
30 appropriate solvent, usually the reaction solvent, for example propan-2-ol. Alternatively, the source of chloride ion may be an aqueous solution of hydrogen chloride, such as concentrated hydrochloric acid.

An alternative source of chloride ion for a salt of Compound (I) is provided by a base salt of hydrochloric acid for example ammonium chloride, or the hydrochloric acid  
35 salt of an amine, for example ethylamine or diethylamine.

The concentration of Compound (I) is preferably in the range of from 3 to 25% weight/volume, more preferably in the range of from 5 to 20%. The concentration of hydrochloric acid solutions are preferably in the range of from 3 to 50% weight/volume.

The reaction is usually carried out at ambient temperature or at an elevated temperature, although any convenient temperature that provides the required product may be employed. A preferred temperature is in the range of from 20-120°C, such as 40°C to 90°C, for example 70°C.

5 Recovery of the required compound generally comprises crystallisation from an appropriate solvent, conveniently the reaction solvent, usually by cooling to a temperature in the range of from 0°C to 40°C, for example 20°C. For example the Hydrochloride may be crystallised from an alcohol such as propan-2-ol, a nitrile such as acetonitrile or an ether such as tetrahydrofuran.

10 In one preferred form the recovery comprises initial crystallisation at an elevated temperature such as 50°C to 90°C preferably 65°C to 75°C and thereafter cooling to a second temperature, suitably in the range of 0°C to 40°C, to complete crystallisation.

Crystallisation can also be initiated by seeding with crystals of the Hydrochloride but this is not essential.

15 Compound (I) is prepared according to known procedures, such as those disclosed in EP 0,306,228 and WO94/05659. The disclosures of EP 0,306,228 and WO94/05659 are incorporated herein by reference.

When used herein the term "T<sub>onset</sub>" is generally determined by Differential Scanning Calorimetry and has a meaning generally understood in the art, as for example  
20 expressed in "Pharmaceutical Thermal Analysis, Techniques and Applications", Ford and Timmins, 1989 as "The temperature corresponding to the intersection of the pre-transition baseline with the extrapolated leading edge of the transition".

When used herein in respect of certain compounds the term "good flow properties" is suitably characterised by the said compound having a Hausner ratio of less  
25 than or equal to 1.5, especially of less than or equal to 1.25.

"Hausner ratio" is an art accepted term. When used herein the term 'prophylaxis of conditions associated with diabetes mellitus' includes the treatment of conditions such as insulin resistance, impaired glucose tolerance, hyperinsulinaemia and gestational diabetes.

30 Diabetes mellitus preferably means Type II diabetes mellitus.

Conditions associated with diabetes include hyperglycaemia and insulin resistance and obesity. Further conditions associated with diabetes include hypertension, cardiovascular disease, especially atherosclerosis, certain eating disorders, in particular the regulation of appetite and food intake in subjects suffering from disorders associated  
35 with under-eating, such as anorexia nervosa, and disorders associated with over-eating, such as obesity and anorexia bulimia. Additional conditions associated with diabetes include polycystic ovarian syndrome and steroid induced insulin resistance.

The complications of conditions associated with diabetes mellitus encompassed herein includes renal disease, especially renal disease associated with the development of Type II diabetes including diabetic nephropathy, glomerulonephritis, glomerular sclerosis, nephrotic syndrome, hypertensive nephrosclerosis and end stage renal disease.

5 As mentioned above the compound of the invention has useful therapeutic properties: The present invention accordingly provides the Hydrochloride for use as an active therapeutic substance.

More particularly, the present invention provides the Hydrochloride for use in the treatment and/or prophylaxis of diabetes mellitus, conditions associated with diabetes  
10 mellitus and certain complications thereof.

Hydrochloride may be administered per se or, preferably, as a pharmaceutical composition also comprising a pharmaceutically acceptable carrier. The formulation of the Hydrochloride is generally as disclosed for Compound (I) in the above mentioned publications.

15 Accordingly, the present invention also provides a pharmaceutical composition comprising the Hydrochloride and a pharmaceutically acceptable carrier therefor.

The Hydrochloride is normally administered in unit dosage form.

The active compound may be administered by any suitable route but usually by the oral or parenteral routes. For such use, the compound will normally be employed in  
20 the form of a pharmaceutical composition in association with a pharmaceutical carrier, diluent and/or excipient, although the exact form of the composition will naturally depend on the mode of administration.

Compositions are prepared by admixture and are suitably adapted for oral, parenteral or topical administration, and as such may be in the form of tablets, capsules,  
25 oral liquid preparations, powders, granules, lozenges, pastilles, reconstitutable powders, injectable and infusable solutions or suspensions, suppositories and transdermal devices. Orally administrable compositions are preferred, in particular shaped oral compositions, since they are more convenient for general use.

Tablets and capsules for oral administration are usually presented in a unit dose,  
30 and contain conventional excipients such as binding agents, fillers, diluents, tableting agents, lubricants, disintegrants, colourants, flavourings, and wetting agents. The tablets may be coated according to well known methods in the art.

Suitable fillers for use include cellulose, mannitol, lactose and other similar agents. Suitable disintegrants include starch, polyvinylpyrrolidone and starch derivatives  
35 such as sodium starch glycolate. Suitable lubricants include, for example, magnesium stearate. Suitable pharmaceutically acceptable wetting agents include sodium lauryl sulphate.

Solid oral compositions may be prepared by conventional methods of blending, filling, tableting or the like. Repeated blending operations may be used to distribute the active agent throughout those compositions employing large quantities of fillers. Such operations are, of course, conventional in the art.

5 Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, gelatin, hydroxyethylcellulose, carboxymethyl  
10 cellulose, aluminium stearate gel or hydrogenated edible fats, emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example, almond oil, fractionated coconut oil, oily esters such as esters of glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl p-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or  
15 colouring agents.

For parenteral administration, fluid unit dose forms are prepared containing a compound of the present invention and a sterile vehicle. The compound, depending on the vehicle and the concentration, can be either suspended or dissolved. Parenteral solutions are normally prepared by dissolving the active compound in a vehicle and filter  
20 sterilising before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents are also dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum.

Parenteral suspensions are prepared in substantially the same manner except that  
25 the active compound is suspended in the vehicle instead of being dissolved and sterilised by exposure to ethylene oxide before suspending in the sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the active compound.

As is common practice, the compositions will usually be accompanied by written  
30 or printed directions for use in the medical treatment concerned.

As used herein the term 'pharmaceutically acceptable' embraces compounds, compositions and ingredients for both human and veterinary use: for example the term 'pharmaceutically acceptable salt' embraces a veterinarily acceptable salt.

The present invention further provides a method for the treatment and/or  
35 prophylaxis of diabetes mellitus, conditions associated with diabetes mellitus and certain complications thereof, in a human or non-human mammal which comprises administering an effective, non-toxic, amount of Hydrochloride to a human or non-human mammal in need thereof.

The compositions are formulated according to conventional methods, such as those disclosed in standard reference texts, for example the British and US Pharmacopoeias, Remington's Pharmaceutical Sciences (Mack Publishing Co.), Martindale The Complete Drug Reference (London, The Pharmaceutical Press) and  
5 Harry's Cosmeticology (Leonard Hill Books). Conveniently, the active ingredient may be administered as a pharmaceutical composition hereinbefore defined, and this forms a particular aspect of the present invention.

In a further aspect the present invention provides the use of Hydrochloride for the manufacture of a medicament for the treatment and/or prophylaxis of diabetes mellitus,  
10 conditions associated with diabetes mellitus and certain complications thereof.

In the treatment and/or prophylaxis of diabetes mellitus, conditions associated with diabetes mellitus and certain complications thereof the Hydrochloride may be taken in amounts so as to provide Compound I in suitable doses, such as those disclosed in EP 0,306,228 and WO94/05659.

15 No adverse toxicological effects are indicated in the above mentioned treatments for the compounds of the invention.

The following examples illustrate the invention but do not limit it in any way.

**Example 1 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione hydrochloride**

A mixture of 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione (9.0 g) and propan-2-ol (180 ml) was stirred and heated to 75°C. A solution of hydrogen chloride in propan-2-ol (7.5 ml, 5-6N) was added and the mixture was stirred for 1 hour at 75°C and then cooled to 21°C. The product was collected by filtration, washed with propan-2-ol (25 ml) and dried under vacuum over phosphorus pentoxide for 16 hours to give the title compound (9.8 g) as a white crystalline solid.

10 Ionic chlorine content: 8.7% wt/wt  
Water content (Karl-Fisher) : 0.3 % wt/wt

<sup>1</sup>H-NMR (d<sup>6</sup>-DMSO): consistent with 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione hydrochloride containing propan-2-ol 2.1% wt/wt.

**Example 2 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione hydrochloride**

20 A mixture of 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione (6.0 g) and propan-2-ol (120 ml) was heated to 50°C before a solution of hydrogen chloride in propan-2-ol (5.0 ml, 5-6 N) was added. The reaction temperature was raised to 70°C and stirring continued at this temperature for 2 hours, during which time crystallisation was observed to have started. After cooling to 21°C the product was collected by filtration, washed with propan-2-ol (20 ml) and dried under vacuum, over phosphorus pentoxide, to give 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione hydrochloride (6.31 g) as a white crystalline solid.

**Example 3 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione hydrochloride**

30 A mixture of 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione (6.0 g) and acetonitrile (120 ml) was stirred and heated to 50°C and a solution of hydrogen chloride in propan-2-ol (5.0 ml, 5-6N) was added. The reaction mixture was warmed to 70°C and stirred at this temperature for 60 minutes, during which time crystallisation was observed to have started. After cooling to 21°C the product was collected by filtration, washed with acetonitrile (20 ml) and then dried under vacuum over phosphorus pentoxide to give 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione hydrochloride (5.7 g) as a white crystalline solid.

**Example 4 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione hydrochloride**

45 A mixture of 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione (6.0 g) and tetrahydrofuran (120 ml) was stirred and heated to 65°C. A solution

of hydrogen chloride in propan-2-ol (5.0 ml, 5-6N) was added and the reaction mixture stirred at 65°C for 2 hours and then cooled to 21°C. The product was collected by filtration, washed with tetrahydrofuran (20 ml) and dried under vacuum over phosphorus pentoxide to give 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione hydrochloride (6.25 g) as white crystalline solid.

**Example 5 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione hydrochloride**

Concentrated hydrochloric acid (0.85ml) was added to a stirred solution of 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione (3.0 g) in propan-2-ol (90 ml) at reflux. The solution was maintained at reflux for 15 minutes and then cooled to 0°C where the precipitation of an oil was observed. The mixture was then warmed to 55°C to produce a clear solution before being cooled to 50°C to initiate crystallisation. The mixture was further cooled to 21°C and the solid was collected by filtration to afford 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione hydrochloride (2.61 g) as white crystalline solid.

**Characterising data recorded for the product of Example 1.**

The infrared absorption spectrum of a mineral oil dispersion of the product was obtained using a Nicolet 710 FT-IR spectrometer at 2 cm<sup>-1</sup> resolution (Figure 1). Data were digitised at 1 cm<sup>-1</sup> intervals. Bands were observed at: 2748, 1742, 1697, 1640, 1604, 1543, 1511, 1413, 1351, 1329, 1251, 1232, 1205, 1179, 1159, 1108, 1057, 1028, 996, 973, 925, 902, 817, 764, 737, 710, 662, 617, 601, 591, 557, 525, 505 cm<sup>-1</sup>.

The infrared spectrum of the solid product was recorded using a Perkin-Elmer Spectrum One FT-IR spectrometer fitted with a universal ATR accessory. Bands were observed at: 2748, 1742, 1687, 1639, 1603, 1543, 1510, 1461, 1413, 1328, 1301, 1250, 1231, 1205, 1179, 1156, 1056, 1028, 996, 973, 925, 901, 817, 762, 738, 709, 661 cm<sup>-1</sup>.

The Raman spectrum of the product (Figure 2) was recorded with the sample in a glass vial using a Perkin-Elmer 2000R FT-Raman spectrometer, at 4 cm<sup>-1</sup> resolution with excitation was from a Nd:YAG laser (1064 nm) with a power output of 400 mW. Bands were observed at: 3100, 3056, 2931, 1744, 1610, 1586, 1544, 1461, 1440, 1377, 1351, 1319, 1289, 1253, 1232, 1208, 1177, 1095, 1028, 998, 980, 925, 900, 840, 825, 772, 740, 714, 664, 637, 619, 606, 469, 430, 397, 340, 300 cm<sup>-1</sup>.

The X-Ray Powder Diffractogram pattern of the product (Figure 3) was recorded using the following acquisition conditions: Tube anode: Cu, Generator tension: 40 kV, Generator current: 40 mA, Start angle: 2.0 °2θ, End angle: 35.0 °2θ, Step size: 0.02 °2θ, Time per step: 2.5 seconds. Characteristic XRPD angles and relative intensities are recorded in Table 1.

Table 1.

Angle 2-Theta °	Rel. Intensity %
5.5	15.4
6.7	14.1
8.7	3.3
11.1	39.4
12.0	9.2
12.7	20.3
13.5	40.8
14.7	40.8
15.1	14.5
15.5	38.1
16.3	30.2
16.8	18.5
17.4	27.1
17.7	34.6
18.2	26.7
18.5	26
18.8	38.8
19.8	31
20.3	11.7
21.2	16.7
21.7	26.4
21.9	26
22.5	14.7
23.2	26.2
23.6	90.8
24.2	100
25.1	49.6
25.5	68.5
25.9	29.3
26.4	16.8
27.3	25.6
27.8	15.4
28.1	28.8
28.8	21.6
29.3	44.1
29.9	20.9
30.5	17.8
30.9	16.8

31.8	24
32.9	18.7
33.1	22
34.3	22.7

The solid-state NMR spectrum of the product (Figure 4) was recorded on a Bruker AMX36 instrument operating at 90.55 MHz: The solid was packed into a 4 mm zirconia MAS rotor fitted with a Kel-F cap and rotor spun at ca.10 kHz. The  $^{13}\text{C}$  MAS spectrum was acquired by cross-polarisation from Hartmann-Hahn matched protons (CP contact time 3 ms, repetition time 15 s) and protons were decoupled during acquisition using a two-pulse phase modulation (TPPM) composite sequence. Chemical shifts were externally referenced to the carboxylate signal of glycine at 176.4 ppm relative to TMS and were observed at: 39.0, 52.5, 54.9, 64.1, 67.3, 111.6, 114.4, 131.0, 137.0, 144.2, 151.3, 156.4, 158.5, 171.6, 175.5, 177.9 ppm.

#### Properties of the Hydrochloride, recorded for the product of Example 1

##### Solid State Stability of the Hydrochloride

The solid state stability of the drug substance was determined by storing approximately 1.0 g of the material in a glass bottle at a) 40°C / 75% Relative Humidity (RH), open exposure, for 1 month and b) at 50°C, closed, for 1 month. The material was assayed by HPLC for final content and degradation products in both cases.

a) 40°C / 75% RH: No significant degradation observed (HPLC assay 100% initial).

b) 50°C: No significant degradation observed (HPLC assay 101% initial).

##### Solubility of the Hydrochloride

The solubility of the material was determined by adding water in aliquots from 1 to 1000ml to approximately 100mg of drug substance until the powder had dissolved. The visual solubility was confirmed by an HPLC assay of a saturated solution.

Solubility: 25 mg/ml.

##### Melting Point of the Hydrochloride

The melting point of the Hydrochloride was determined according to the method described in the U.S. Pharmacopoeia, USP 23, 1995, <741> "Melting range or temperature, Procedure for Class Ia", using a Buchi 545 melting point instrument.

Melting Point: 167°C

##### T<sub>onset</sub> of the Hydrochloride

The T<sub>onset</sub> of the drug substance was determined by Differential Scanning Calorimetry using a Perkin-Elmer DSC apparatus.

Product of example 1: T<sub>onset</sub> (10°C/minute, open pan): 165°C


Product of example 2: T<sub>onset</sub> (10°C/minute, open pan): 163°C

## CLAIMS:


1. A compound 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione, hydrochloride salt characterised in that it provides:
  - 5 (i) an infrared spectrum substantially in accordance with Figure 1; and/or
  - (ii) a Raman spectrum substantially in accordance with Figure 2; and/or
  - (iii) an X-Ray powder diffraction pattern (XRPD) substantially in accordance with Figure 3; and
  - 10 (iv) a Solid State  $^{13}\text{C}$  NMR spectrum substantially in accordance with Figure 4.
2. A compound according to claim 1, characterised in that it provides two or more of:
  - (i) provides an infrared spectrum substantially in accordance with Figure 1;
  - (ii) a Raman spectrum substantially in accordance with Figure 2;
  - 15 (iii) an X-Ray powder diffraction pattern (XRPD) substantially in accordance with Figure 3;
  - (iv) a Solid State  $^{13}\text{C}$  NMR spectrum substantially in accordance with Figure 4; and
  - (v) a melting point in the range of from 160 to 168°C.
- 20 3. A compound according to claim 1 or claim 2, in purified form.
4. A compound according to claim 1 or claim 2, in a solid dosage form.
5. A compound according to claim 1 or claim 2, in a pharmaceutically acceptable  
25 form capable of being milled or in a milled form
6. A compound according to claim 1 or claim 2, in a pharmaceutically acceptable form and having good flow properties.
- 30 7. A process for preparing the Hydrochloride, according to claim 1, characterised in that 5-[4-[2-(N-methyl-N-(2-pyridyl)amino)ethoxy]benzyl]thiazolidine-2,4-dione (Compound I) or salt thereof, preferably dispersed or dissolved in a suitable solvent, is reacted with a source of chloride ion; and thereafter the Hydrochloride is recovered.
- 35 8. A pharmaceutical composition comprising the Hydrochloride according to claim 1, and a pharmaceutically acceptable carrier therefor.
9. A Hydrochloride, according to claim 1, for use as an active therapeutic substance.

10. A use of the Hydrochloride according to claim 1, for the manufacture of a medicament for the treatment and/or prophylaxis of diabetes mellitus, conditions associated with diabetes mellitus and certain complications thereof.


1/4

**Fig.1** Infrared spectrum of the Hydrochloride


2/4

**Fig. 2** Raman spectrum of the Hydrochloride

3/4

**Fig. 3** X-Ray Powder Diffractogram of the Hydrochloride

4/4

**Fig. 4** Solid State  $^{13}\text{C}$  NMR spectrum of the Hydrochloride

## INTERNATIONAL SEARCH REPORT

International Application No

PC1/GB 01/03991

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D417/12 A61K31/44 A61P3/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94 05659 A (SMITHKLINE BEECHAM PLC ;POOL COLIN RIPLEY (GB); ROMAN ROBIN SHERWO) 17 March 1994 (1994-03-17) cited in the application page 2, line 13 - line 15; example 1 -----	1-10
X	EP 0 306 228 A (BEECHAM GROUP PLC) 8 March 1989 (1989-03-08) cited in the application claims 1,12; example 30 -----	1-10

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the International filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&amp;\* document member of the same patent family

Date of the actual completion of the international search

3 December 2001

Date of mailing of the international search report

10/12/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Fritz, M

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/03991

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9405659	A	17-03-1994	AP 513 A	30-07-1996
			AT 182147 T	15-07-1999
			AU 674880 B2	16-01-1997
			AU 4973093 A	29-03-1994
			BR 1100916 A3	04-07-2000
			CA 2143849 A1	17-03-1994
			CN 1101911 A ,B	26-04-1995
			CN 1183275 A ,B	03-06-1998
			CN 1183413 A ,B	03-06-1998
			CN 1183276 A ,B	03-06-1998
			CZ 9500565 A3	15-11-1995
			DE 69325658 D1	19-08-1999
			DE 69325658 T2	30-12-1999
			DK 658161 T3	29-11-1999
			EP 0658161 A1	21-06-1995
			EP 0960883 A1	01-12-1999
			ES 2133410 T3	16-09-1999
			FI 951004 A	03-03-1995
			FI 982413 A	06-11-1998
			WO 9405659 A1	17-03-1994
			GR 3030794 T3	30-11-1999
			HK 1012363 A1	05-05-2000
			HU 72639 A2	28-05-1996
			IL 106904 A	30-09-1997
			JP 11147885 A	02-06-1999
			JP 2828777 B2	25-11-1998
			JP 8501095 T	06-02-1996
			LU 90712 A9	12-03-2001
			MX 9305397 A1	31-01-1995
			NO 950852 A	03-03-1995
			NO 974646 A	03-03-1995
			NZ 255505 A	22-08-1997
			PL 307812 A1	26-06-1995
			RU 2128179 C1	27-03-1999
			SG 48302 A1	17-04-1998
			SI 9300452 A	30-06-1994
			SK 27795 A3	09-08-1995
			TW 385309 B	21-03-2000
			US 5741803 A	21-04-1998
			US 5910592 A	08-06-1999
			ZA 9306509 A	16-06-1994
EP 0306228	A	08-03-1989	AT 186724 T	15-12-1999
			AU 2173888 A	09-03-1989
			BR 1100841 A3	20-06-2000
			CA 1328452 A1	12-04-1994
			CA 1339902 A1	09-06-1998
			CZ 9103916 A3	17-03-1993
			DE 3856378 D1	23-12-1999
			DE 3856378 T2	11-05-2000
			DK 490288 A	05-03-1989
			DK 200001556 A	18-10-2000
			EP 0306228 A1	08-03-1989
			EP 0842925 A1	20-05-1998
			ES 2137915 T3	01-01-2000
			GR 3031873 T3	29-02-2000
			HK 1011029 A1	03-11-2000
			JP 10194970 A	28-07-1998

## INTERNATIONAL SEARCH REPORT

In tional Application No

PCT/GB 01/03991

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0306228 A		JP 10194971 A	28-07-1998
		JP 1131169 A	24-05-1989
		JP 2614497 B2	28-05-1997
		JP 2817840 B2	30-10-1998
		JP 9183771 A	15-07-1997
		JP 2837139 B2	14-12-1998
		JP 9183726 A	15-07-1997
		JP 9183772 A	15-07-1997
		KR 164207 B1	15-01-1999
		KR 164275 B1	15-01-1999
		KR 169463 B1	15-01-1999
		LU 90711 A9	05-03-2001
		NZ 226027 A	26-03-1992
		PT 88410 A ,B	31-07-1989
		SG 59988 A1	22-02-1999
		SK 391691 A3	11-12-2000
		US 6288095 B1	11-09-2001
		US 5646169 A	08-07-1997
		US 5002953 A	26-03-1991
		US 5521201 A	28-05-1996
		US 5232925 A	03-08-1993
		US 5194443 A	16-03-1993
		US 5756525 A	26-05-1998
		US 5260445 A	09-11-1993
		ZA 8806536 A	26-07-1989